

H3C1 Polyclonal Antibody

Catalog Number:D-AB-10286L

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human,Mouse,Rat
Immunogen	Recombinant Human H3C1 protein expressed by E.coli
Host	Rabbit
Isotype	IgG
Purification	Antigen Affinity Purification
Conjugation	Unconjugated
Formulation	PBS with 0.02% sodium azide,50% glycerol pH 7.4

Applications Recommended Dilution

WB	1:500-1:1000
-----------	--------------

Data



Western blot with H3C1 Polyclonal antibody at dilution of 1:1000.lane 1:C6 whole cell lysate,lane 2:NIH/3T3 whole cell lysate,lane 3:Jurkat whole cell lysate,lane 4:Hela whole cell lysate.

Observed Mw:15kDa
Calculated Mw:15kDa

Preparation & Storage

Storage Store at -20°C. Avoid freeze / thaw cycles.

Background

Histones are small,highly basic proteins that consist of a globular domain with unstructured N- and C-terminal tails protruding from the main structure. Histone H3 is one of the five main histones that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A,H2B,H3,and H4) form an octamer,around which approximately 146 bp of DNA is wrapped in repeating units,called nucleosomes. In addition to their role in DNA compartmentalization,histones also play crucial roles in various biologic processes,including gene expression and regulation,DNA repair,chromatin condensation,cell cycle progression,chromosome segregation,and apoptosis. The ability of histones to regulate chromatin dynamics primarily originates from various posttranslational modifications carried out by histone-modifying enzymes.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017