

PE Anti-Human CD99 Antibody[HI156]

Catalog No.	E-AB-F1339D	Reactivity	Human
Storage	Store at 2~8°C, Avoid freeze / thaw cycles	Applications	FCM

Important Note: Centrifuge before opening to ensure complete recovery of vial contents.

Antigen Information

Alternate Names	CD99, MIC2, HBA71, MSK5X, E2 antigen
Uniprot ID	P14209
Gene ID	4267
Background	CD99 is a type I single chain transmembrane protein devoid of N-linked glycosylation sites encoded by the pseudoautosomal gene MIC2. CD99 has an apparent molecular weight of 32 kD and is widely expressed on a variety of tissues. CD99 is highly expressed on thymocytes, T cells, and T cell leukemias and lymphomas. However, it is absent on some B cell lines, fetal B cells, eosinophils, granulocytes and the NK-cell line YT. CD99 is involved in spontaneous rosette formation with erythrocytes and may also be involved in other T-cell and hematopoietic cell adhesion pathways. CD99 has been reported to activate a caspase-independent death pathway in T cells under some conditions. CD99 interacts with a number of proteins including ferritin heavy chain 1, karyopherin beta 1, TRIP13, cyclophilin A, annexin II, and ubiquitin-conjugating enzyme E2H.

Product Details

Form	Liquid
Size	20Tests/100Tests/100Tests×2
Clone No.	HI156
Host	Mouse
Isotype	Mouse IgG2a, κ
Reactivity	Human
Application	FCM
Isotype Control	PE Mouse IgG2a, κ Isotype Control[C1.18.4] [Product E-AB-F09802D]
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.
Shipping	Biological ice pack at 4 °C
Stability & Storage	Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light.Do not freeze. This product is guaranteed up to one year from purchase.

For Research Use Only

Fluorophore


Conjugation: PE

PE is designed to be excited by the Blue (488 nm), Green (532 nm) and Yellow-Green (561 nm) lasers and detected using an optical filter centered near 575 nm (e.g., a 585/42 nm bandpass filter).

Recommended usage

Each lot of this antibody is quality control tested by flow cytometric analysis. **The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood).** Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Product data



Human peripheral blood lymphocytes are stained with PE Anti-Human CD99 Antibody[HI156] (filled gray histogram) or PE Mouse IgG2a, κ Isotype Control (empty black histogram).

Related Information

1. Sample Preparation for Flow Cytometry <https://www.elabscience.com/List-detail-5594.html>
2. Staining Cell Surface Targets for Flow Cytometry <https://www.elabscience.com/List-detail-5568.html>
3. Flow Cytometry Troubleshooting Tips <https://www.elabscience.com/List-detail-5593.html>
4. How to select the appropriate detection channel through the spectrogram? <https://www.elabscience.com/List-detail-459742.html>